



Some graft transformations and its applications on the distance spectral radius of a graph

Guanglong Yu^{a,b}, Huicai Jia^d, Hailiang Zhang^{c,b}, Jinlong Shu^{b,*}

^a Department of Mathematics, Yancheng Teachers University, Yancheng, 224002, Jiangsu, China

^b Department of Mathematics, East China Normal University, Shanghai, 200241, China

^c Department of Mathematics, Taizhou University, Taizhou, 317000, Zhejiang, China

^d Department of Mathematical and Physical Sciences, Henan Institute of Engineering, Zhengzhou, Henan, 451191, China

ARTICLE INFO

Article history:

Received 24 March 2011

Received in revised form 6 September 2011

Accepted 7 September 2011

Keywords:

Graft transformation

Distance spectral radius

Pendant number

ABSTRACT

Let $D(G) = (d_{ij})_{n \times n}$ denote the distance matrix of a connected graph G with order n , where d_{ij} is equal to the distance between v_i and v_j in G . The largest eigenvalue of $D(G)$ is called the distance spectral radius of graph G , denoted by $\varrho(G)$. In this paper, some graft transformations that decrease or increase $\varrho(G)$ are given. With them, for the graphs with both order n and k pendant vertices, the extremal graphs with the minimum distance spectral radius are completely characterized; the extremal graph with the maximum distance spectral radius is shown to be a dumbbell graph (obtained by attaching some pendant edges to each pendant vertex of a path respectively) when $2 \leq k \leq n - 2$; for $k = 1, 2, 3, n - 1$, the extremal graphs with the maximum distance spectral radius are completely characterized.

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

The distance matrix of a graph, while not as common as the more familiar adjacency matrix, has nevertheless come up in several different areas, including communication network design [1], graph embedding theory [2–4], molecular stability [5,6] and network flow algorithms [7,8]. So it is very interesting to study the spectra of these matrices.

Throughout this article, all graphs considered are simple, connected and undirected. Let $G = G[V(G), E(G)]$ be a graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G)$, where $|V(G)| = n$ is the order and $|E(G)| = m$ is the size of G . Let $N_G(v)$ denote the set of the vertices adjacent to v in G . The degree of v in G , denoted by $\deg(v)$, is equal to $|N_G(v)|$. A vertex is called a pendant vertex if its degree is 1. We denote by P_n, K_n, S_n a path, a complete graph, a star with order n , respectively. In a graph G , the length of a shortest path from v_i to v_j is called the distance between v_i and v_j , denoted by $d(v_i, v_j)$ or $d_G(v_i, v_j)$. $d(G) = \max d(v_i, v_j) \mid v_i, v_j \in V(G)$ is called the *diameter* of graph G . Let $P_{i,j}$ denote a shortest path from vertex v_i and v_j . If the length of the path $P_{i,j}$ is equal to $d(G)$, then $P_{i,j}$ is called a *diameter-path*, and v_i, v_j are called the end vertices of $P_{i,j}$. For $S \subseteq V(G)$, let $G[S]$ denote the subgraph induced by S . For a vertex set $\{v_1, v_2, \dots, v_k\}$, we sometimes abbreviate $G[\{v_1, v_2, \dots, v_k\}]$ as $G[v_1, v_2, \dots, v_k]$. The dumbbell graph $D(n; s, p)$ consists of a path P_{n-s-p} , s pendant edges attaching to a pendant vertex of P_{n-s-p} and p pendant edges attaching to the other pendant vertex of P_{n-s-p} ($s \geq 0, p \geq 0, s + p \leq n - 2$). Denote by $\mathcal{D}(n, k; s, p) = \{D(n; s, p) \mid s + p = k\}$.

Let $D(G) = (d_{ij})_{n \times n}$ denote the distance matrix of a connected graph G with order n , where $d_{ij} = d_G(v_i, v_j)$. The distance characteristic polynomial of G , denoted by $P(D(G))$ (or $P(D(G), \lambda)$), is defined as $\det(\lambda I - D(G))$, where I is the

* Corresponding author.

E-mail addresses: yglong01@163.com (G. Yu), jhc607@yahoo.com.cn (H. Jia), rockzhang76@tzc.edu.cn (H. Zhang), jlshu@math.ecnu.edu.cn (J. Shu).





Fig. 1.1. Graft transformation.

Fig. 2.1. G, H .

identity matrix. The largest eigenvalue of $P(D(G))$ is called the distance spectral radius of G , denoted by $\varrho(G)$. By the Perron–Frobenius theorem [9], for a connected graph G , we know that there exists a unit positive vector corresponding to $\varrho(G)$ which is called *Perron eigenvector*. The pendant number of a graph G , denoted by $\mathcal{K}(G)$, is the number of pendant vertices. Let $\mathcal{G}(n; k) = \{G \mid G \text{ be a graph with order } n, \mathcal{K}(G) = k\}$, $\mathcal{T}(n; k) = \{T \mid T \text{ be a tree with order } n \text{ and } \mathcal{K}(T) = k\}$, $\varrho_{\max}(\mathcal{G}; n, k) = \max\{\varrho(G) \mid G \in \mathcal{G}(n; k)\}$, $\varrho_{\min}(\mathcal{G}; n, k) = \min\{\varrho(G) \mid G \in \mathcal{G}(n; k)\}$, $\varrho_{\max}(\mathcal{T}; n, k) = \max\{\varrho(T) \mid T \in \mathcal{T}(n; k)\}$, $\varrho_{\min}(\mathcal{T}; n, k) = \min\{\varrho(T) \mid T \in \mathcal{T}(n; k)\}$.

Let u, v be two vertices of a connected graph G . Suppose $v_1, v_2, \dots, v_s (1 \leq s \leq \deg(v))$ are some vertices of $N_G(v) \setminus N_G[u] (N_G[u] = N_G(u) \cup \{u\})$. Let G^* be the graph obtained from G by deleting the edges (v, v_i) and adding the edges $(u, v_i) (1 \leq i \leq s)$ (see Fig. 1.1). We call the process from G to G^* a *graft transformation*. An interesting thing is that a graft transformation always changes the distance spectral radius (adjacency spectral radius [10], signless Laplacian spectral radius [11]) of a graph. In this paper, some graft transformations that decrease or increase $\varrho(G)$ are presented. With them, for the graphs with both order n and k pendant vertices, the extremal graphs with the minimum distance spectral radius are completely characterized; the extremal graph with the maximum distance spectral radius is shown to be a dumbbell graph when $2 \leq k \leq n - 2$; for $k = 1, 2, 3, n - 1$, the extremal graphs with the maximum distance spectral radius are completely characterized respectively.

2. Graft transformations

Lemma 2.1 ([12]). Let A be an $n \times n$ real symmetric irreducible nonnegative matrix and $X \in \mathbb{R}^n$ be a unit vector. If $\rho(A) = X^T A X$, then $A X = \rho(A) X$.

Definition 2.2. The union of simple graphs H and G is the simple graph $G \cup H$ with vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H)$. The intersection $G \cap H$ of simple graphs H and G is defined analogously.

Theorem 2.3. Suppose graph $G = \bigcup_{i=1}^3 G_i$ satisfies that $G_i \cap G_j = v_0$ for $1 \leq i, j \leq 3, i \neq j$, and that $|V(G_i)| \geq 2$ for $i = 1, 2, 3$ (see Fig. 2.1). $X = (x_0, x_1, \dots, x_{n-1})^T$ is the Perron eigenvector corresponding to $\varrho(G)$, in which x_i corresponds to v_i . Let $S_1 = \sum_{v_i \in V(G_1)} x_i$, $S_2 = \sum_{v_i \in V(G_2)} x_i$, and for a vertex $v_a \in V(G_2)$, $v_a \neq v_0$, let graph $H = G - \sum_{v_i \in N_{G_3}(v_0)} v_0 v_i + \sum_{v_i \in N_{G_3}(v_0)} v_a v_i$. If $S_1 \geq S_2$, then $\varrho(H) > \varrho(G)$.



Proof. Suppose that $P_{a,0}$ is a shortest path from v_a to v_0 in G_2 with length $d_a \geq 1$. Note that for $v_i \in V(G_3) \setminus \{v_0\}$, $v_j \in V(G_1)$, we have $d_H(v_i, v_j) - d_G(v_i, v_j) = d_a$; for $v_i \in V(G_3) \setminus \{v_0\}$, $v_k \in V(G_2)$, we have $d_G(v_i, v_k) - d_H(v_i, v_k) \leq d_a$. So

$$\varrho(H) - \varrho(G) \geq X^T D(H) X - X^T D(G) X \geq 2d_a(S_1 - S_2) \sum_{v_i \in V(G_3) \setminus \{v_0\}} x_i \geq 0.$$

Suppose $\varrho(H) = \varrho(G)$, then $X^T D(H) X = \varrho(H)$. By Lemma 2.1, $D(H) X = \varrho(H) X$, but $\varrho(H) x_0 = D(H)_0 X > D(G)_0 X = \varrho(G) x_0$, where $D(H)_0, D(G)_0$ denote the rows corresponding to v_0 in $D(H), D(G)$ respectively. So, $\varrho(H) \neq \varrho(G)$ and then $\varrho(H) > \varrho(G)$. \square

Same as proof of Theorem 2.3, we can prove the following theorem.

Theorem 2.4. Graph G as shown in Fig. 2.2 satisfies that $G_1 \cap G_3 = v_1$, $v_1 v_2$ is a cut edge. $X = (x_1, x_2, \dots, x_n)^T$ is the Perron eigenvector corresponding to $\varrho(G)$, in which x_i corresponds to v_i . Let $S_1 = \sum_{v_i \in V(G_1)} x_i$, $S_2 = \sum_{v_i \in V(G_2)} x_i$, and let graph $H = G - \sum_{v_i \in N_{G_3}(v_1)} v_1 v_i + \sum_{v_i \in N_{G_3}(v_1)} v_2 v_i$. If $S_1 \geq S_2$, $|V(G_3)| \geq 2$, then $\varrho(H) > \varrho(G)$.

Fig. 2.2. G, H .Fig. 3.1. G .Fig. 3.2. H .

3. Extremal graphs

For a nonnegative irreducible square matrix A , the *spectral radius*, denoted by $\rho(A)$, is the maximum of the moduli of its eigenvalues.

Lemma 3.1 ([13]). Suppose both A, B are nonnegative irreducible and $B \leq A$ (namely $B_{ij} \leq A_{ij}$ for each pair of i, j). Then $\rho(B) \leq \rho(A)$ with equality if and only if $B = A$.

Lemma 3.2 ([14]). Suppose that G_1 is a complete graph with $V(G_1) = \{v_0, v_{k+1}, v_{k+2}, \dots, v_{n-1}\}$ ($n-3 \geq k$). Graph G consists of the complete graph G_1 and pendant edges $v_0v_1, v_0v_2, \dots, v_0v_k$. Graph H consists of G_1 and pendant stars S_{v_i} attached at each vertex v_i (v_i is the center of S_{v_i}) of the complete graph G_1 where stars can be trivial (with only one vertex). Then we have

- (i) if $k = 0, 1$, then $\rho(H) = \rho(G)$;
- (ii) if $k \geq 2$ and $2 \leq t_0 < k+1$, then $\rho(H) > \rho(G)$.

Theorem 3.3. If graph $G \in \mathcal{G}(n; k)$ satisfies that $\rho(G) = \rho_{\min}(\mathcal{G}; n, k)$, then

- (i) for $n = 2, G \cong K_2$;
- (ii) for $n \geq 1, n \neq 2, 0 \leq k \leq n-1, k \neq n-2$, G is isomorphic to a graph obtained by attaching k pendant edges to a vertex of a complete graph with order $n-k$;
- (iii) for $n \geq 4, k = n-2, G \cong D(n; 1, n-3)$.

Proof. (i) clearly holds.

(ii) clearly holds for $k = n-1$. Next we consider the case that $k \leq n-3$. Suppose that v_1, v_2, \dots, v_{n-k} are the non-pendant vertices of G . By Lemma 3.1, we know that for a connected graph \mathcal{G} , if $uv \notin E(\mathcal{G})$ ($u, v \in V(\mathcal{G})$), then $\rho(\mathcal{G} + uv) < \rho(\mathcal{G})$. Therefore, in G , if $G[v_1, v_2, \dots, v_{n-k}]$ is not a clique (complete subgraph), by adding edges to $G[v_1, v_2, \dots, v_{n-k}]$, we can get a new graph H satisfying that $H[v_1, v_2, \dots, v_{n-k}]$ is a clique. Then $\rho(H) < \rho(G)$, which contradicts $\rho(G) = \rho_{\min}(\mathcal{G}; n, k)$. So $G[v_1, v_2, \dots, v_{n-k}]$ must be a clique. Then for $k \leq n-3$, (ii) follows from Lemma 3.2.


It is easy to check that (iii) holds for $n = 4, 5$ (see Fig. 3.1). Next we suppose $n \geq 6$. $H \cong D(n; 1, n-3)$ is as shown in Fig. 3.2. $X = (x_1, x_2, \dots, x_n)^T$ is the Perron eigenvector corresponding to $\rho(H)$, in which x_i corresponds to v_i .

By symmetry, $x_4 = x_5 = \dots = x_n$. Let $M = \sum_{i=3}^n x_i$. Note that $\rho(H)x_1 - \rho(H)x_2 = M + x_2 - x_1$. Then $(\rho(H) + 1)(x_1 - x_2) = M$. So we have $x_1 > x_2$. Note that $\rho(H)x_1 - \rho(H)x_4 = (n-1)x_4 + x_3 - x_2 - 3x_1$. So $(\rho(H) + n-1)(x_1 - x_4) = (n-4)x_1 + x_3 - x_2$. Then we have $x_1 > x_4$. Note that $\rho(H)(x_3 + x_4 + x_5) - \rho(H)(x_1 + x_2) = 4x_2 + 7x_1 - 4x_4 - x_3$. So we have $(\rho(H) + 1)(x_3 + x_4 + x_5 - (x_1 + x_2)) = 3x_2 + 6x_1 - 2x_4 > 0$. Then $x_3 + x_4 + x_5 > x_1 + x_2$, and so $x_3 + x_{n-1} + x_n > x_1 + x_2$. For $4 \leq j \leq n-1$, we let $H' = H - \sum_{i=4}^j v_3v_i + \sum_{i=4}^j v_2v_i$. When $j = n-1$, Then $H' \cong H$. When $4 \leq j \leq n-2$, by Lemma 2.4, we get $\rho(H') > \rho(H)$. With these, we get that for $k = n-2$, if $\rho(G) = \rho_{\min}(\mathcal{G}; n, k)$, then $G \cong H$. Then (iii) follows. \square

Theorem 3.4. Suppose that graph $G \in \mathcal{T}(n; k)$ and $\rho(G) = \rho_{\max}(\mathcal{T}; n, k)$ ($2 \leq k \leq n-2$). Then $G \in \mathcal{D}(n, k; s, p)$.

Proof. Assume that the theorem does not hold. Suppose that $P_d = P_1 \cup P_2$ is a diameter-path of G , where path $P_1 = v_s v_{s-1} \dots v_1 v_0$, $P_2 = v_0 v_{s+1} v_{s+2} \dots v_d$. Suppose $G = \bigcup_{i=1}^3 T_i$, where $T_i \cap T_j = v_0$ for $j \neq i, 1 \leq i, j \leq 3$, and $P_1 \subseteq T_1, P_2 \subseteq T_2$. If $d(T_3) \geq 2$, then both the lengths of P_1 and P_2 are at least 2. Suppose that $X = (x_0, x_1, \dots, x_{n-1})^T$ is the Perron eigenvector corresponding to $\rho(G)$, in which x_i corresponds to v_i . Let $S_1 = \sum_{v_i \in V(T_1)} x_i, S_2 = \sum_{v_i \in V(T_2)} x_i$. Suppose $S_1 \geq S_2$. Let $H = G - \sum_{v_i \in N_{T_3}(v_0)} v_0v_i + \sum_{v_i \in N_{T_3}(v_0)} v_{d-1}v_i$. Then $\rho(H) > \rho(G)$ by Theorem 2.3 and $d(H) \geq d(G) + 1$. Proceeding like this, we get a caterpillar graph \mathcal{H} with diameter $n-k+1$ (see Fig. 3.3).

Suppose that $P = v_{a(1,1)} v_1 v_2 \dots v_{n-k-1} v_{n-k} v_{a(n-k,1)}$ is a diameter-path in \mathcal{H} . Denote by $v_i v_{a(i,j)}$ ($0 \leq j \leq t_i$) the pendant edges attaching to vertex v_i ($1 \leq i \leq n-k$). If there exists $t_b > 0, 2 \leq b \leq n-k-1$, let $S_b = \bigcup_{j=1}^{t_b} v_b v_{a(b,j)}$. Suppose that

Fig. 3.3. \mathcal{H} .

$\mathcal{H} = T'_1 \cup T'_2 \cup \mathbb{S}_b$, $T'_1 \cap T'_2 = T'_1 \cap \mathbb{S}_b = T'_2 \cap \mathbb{S}_b = v_b$, $v_1 \in V(T'_1)$, $v_{n-k} \in V(T'_2)$. Suppose that $Y = (y_0, y_1, \dots, y_{n-1})^T$ is the Perron eigenvector corresponding to $\varrho(\mathcal{H})$, in which y_i corresponds to v_i . Let $S'_1 = \sum_{v_i \in V(T'_1)} y_i$, $S'_2 = \sum_{v_i \in V(T'_2)} y_i$. Suppose $S'_1 \geq S'_2$. Let $\mathcal{H}_1 = \mathcal{H} - \sum_{j=1}^{t_b} v_b v_{a(b,j)} + \sum_{j=1}^{t_b} v_{n-k} v_{a(b,j)}$. Then $\varrho(\mathcal{H}_1) > \varrho(\mathcal{H})$ by Theorem 2.3. Proceeding like this, we get a graph $\mathcal{H} \in \mathcal{D}(n, k; s, p)$ such that $\varrho(\mathcal{H}) > \varrho(\mathcal{H}_1)$, which contradicts $\varrho(G) = \varrho_{\max}(\mathcal{D}; n, k)$. Therefore, the theorem holds. \square

Lemma 3.5 ([15]). Denote by $G(v, k)$ the graph obtained from a graph G by attaching a pendant path $P = v_0 v_1 v_2 \dots v_k$ to a vertex $v = v_0$ of G , and denote by ϱ the distance spectral radius of $G(v, k)$. Let $X = (x_0, x_1, \dots, x_{n-1})^T$ be the Perron vector corresponding to ϱ , in which x_i corresponds to v_i , and let $S = \sum_{i=0}^{n-1} x_i$. Then $\sum_{i=1}^k x_i = x_0 f(k) + \frac{S}{\varrho} g(k)$, where $\varrho = \varrho(G(v, k))$, $f(x) = \frac{t(t^{2x}-1)}{(1+t^{2x+1})(t-1)}$ and $g(x) = \frac{t(t^{x-1}(t^{x+1}-1))}{(1+t^{2x+1})(t-1)^2}$ with $t = 1 + \frac{1}{\varrho} + \frac{\sqrt{2\varrho+1}}{\varrho}$. Furthermore, the functions $f(x)$ and $g(x)$ are monotonically increasing.

Lemma 3.6 ([16]). Let w be a vertex of a nontrivial connected graph G and for nonnegative integers p and q , and let $G(p, q)$ denote the graph obtained from G by attaching pendant paths $P = w v_1 v_2 \dots v_p$ and $Q = w u_1 u_2 \dots u_q$. If $p \geq q \geq 1$, then $\varrho(G(p, q)) < \varrho(G(p+1, q-1))$.

Theorem 3.7. If graph $G \in \mathcal{G}(n; k)$ with $2 \leq k \leq n-2$ and $\varrho(G) = \varrho_{\max}(\mathcal{G}; n, k)$, then

- (i) $G \in \mathcal{D}(n, k; s, p)$.
- (ii) Suppose that G is obtained by attaching pendant edges $v_1 v_{n-k+1}, v_1 v_{n-k+2}, \dots, v_1 v_{n-k+s}$ to vertex v_1 of a path $P = v_1 v_2 \dots v_{n-k-1} v_{n-k}$ and attaching pendant edges $v_{n-k} v_{n-k+s+1}, v_{n-k} v_{n-k+s+2}, \dots, v_{n-k} v_n$ to vertex v_{n-k} . $X = (x_1, x_2, \dots, x_n)^T$ is the Perron eigenvector corresponding to $\varrho(G)$, in which x_i corresponds to v_i . Let $S_1 = \sum_{i=n-k+1}^{n-k+s} x_i$, $S_2 = \sum_{i=n-k+s+1}^n x_i$. If $s \geq p$, then $S_1 \geq S_2$, but $x_{n-k+1} \leq x_{n-k+s+1}$. In particular, $x_{n-k+1} < x_{n-k+s+1}$ if $s > p$.

Proof. (i) By Theorem 3.4, we may assume that G is not a tree. Suppose that H is a spanning tree of G . By Lemma 3.1, we know that for a simple connected graph \mathcal{B} , if $uv \in E(\mathcal{B})$ and $\mathcal{B} - uv$ is also connected, then $\varrho(\mathcal{B} - uv) > \varrho(\mathcal{B})$. So, $\varrho(H) > \varrho(G)$. Suppose $\mathcal{K}(H) = \varepsilon$, then $\varepsilon \geq k$. By Theorem 3.4, we can get a graph $H_1 \in \mathcal{D}(n, \varepsilon; r, t)$ such that $\varrho(H_1) \geq \varrho(H)$. Suppose H_1 is obtained by attaching r pendant edges to the vertex v_1 of a path $P' = v_1 v_2 \dots v_{n-\varepsilon}$ and attaching t pendant edges to the vertex $v_{n-\varepsilon}$.

If $\max\{r, t\} \geq k$, for convenience, we suppose $\max\{r, t\} = r$. Using Lemma 3.6 repeatedly, we can get a graph H_2 that $\varrho(H_2) > \varrho(H_1)$ from H_1 by deleting $r - k + 2$ pendant edges attaching at vertex v_1 and deleting all the pendant edges attaching at vertex $v_{n-\varepsilon}$, and then attaching a pendant path with length $r - k + 2$ to vertex v_1 and attaching a pendant path with length t to vertex $v_{n-\varepsilon}$.

If $\max\{r, t\} < k$, for convenience, we suppose $\max\{r, t\} = r$. Using Lemma 3.6 repeatedly, we can get a graph H_2 that $\varrho(H_2) > \varrho(H_1)$ from H_1 by deleting $\varepsilon - k + 1$ pendant edges attaching at vertex $v_{n-\varepsilon}$, and then attaching a pendant path with length $\varepsilon - k + 1$ to vertex $v_{n-\varepsilon}$.

For both above two cases, $\mathcal{K}(H_2) = k$. By Theorem 3.4, we can get a graph $\mathcal{H} \in \mathcal{D}(n, k; s, p)$ such that $\varrho(\mathcal{H}) > \varrho(H_2)$, which contradicts $\varrho(G) = \varrho_{\max}(\mathcal{G}; n, k)$. Therefore, (i) holds.

(ii) Suppose $s \geq p$. By symmetry, $x_{n-k+1} = x_{n-k+2} = \dots = x_{n-k+s}$, $x_{n-k+s+1} = x_{n-k+s+2} = \dots = x_n$. Assume that $S_2 > S_1$, then $x_{n-k+1} < x_{n-k+s+1}$. By Lemma 3.5, we get $x_1 < x_{n-k}$.

1° $n - k$ is odd. Note that, for $1 \leq i \leq \lfloor \frac{n-k}{2} \rfloor - 1$,

$$\varrho(G)(x_{n-k-i} - x_{i+1}) - \varrho(G)(x_{n-k-i+1} - x_i) = 2 \left(\sum_{j=n-k-i+1}^{n-k} x_j + S_2 \right) - 2 \left(\sum_{j=1}^i x_j + S_1 \right) > 0.$$

Because $x_1 < x_{n-k}$, by induction, we can get $x_{n-k-i} - x_{i+1} > 0$ for $0 \leq i \leq \lfloor \frac{n-k}{2} \rfloor - 1$. But

$$\begin{aligned} \varrho(G) \left(x_{\frac{n-k+3}{2}} - x_{\frac{n-k-1}{2}} \right) &= 2 \left(\sum_{j=1}^{\frac{n-k-1}{2}} x_j + S_1 \right) - 2 \left(\sum_{j=\frac{n-k+3}{2}}^{n-k} x_j + S_2 \right) \\ &\Rightarrow (\varrho(G) + 2) \left(x_{\frac{n-k+3}{2}} - x_{\frac{n-k-1}{2}} \right) = 2 \left(\sum_{j=1}^{\frac{n-k-3}{2}} x_j + S_1 \right) - 2 \left(\sum_{j=\frac{n-k+5}{2}}^{n-k} x_j + S_2 \right) < 0, \end{aligned}$$

which contradicts that $x_{n-k-i} - x_{i+1} > 0$ for $0 \leq i \leq \lfloor \frac{n-k}{2} \rfloor - 1$.

2° $n - k$ is even. As 1° , we can get the same contradiction.

From 1° and 2° , we get $S_1 \geq S_2$. Assume that $x_{n-k+1} \geq x_{n-k+s+1}$ if $s > p$, we can get the same contradictions as 1° and 2° . Therefore, (ii) holds. \square

Lemma 3.8 ([14]). Let G be a connected unicyclic graph with order n ($m(G) = n$, $n \geq 3$). Then $\varrho(G) \leq \varrho(P'_n)$ where P'_n is obtained from a triangle K_3 by attaching pendant path with length $n - 3$ to one of its vertices, with equality if and only if $G \cong P'_n$.

Theorem 3.9. If graph $G \in \mathcal{G}(n; k)$ with n and $\varrho(G) = \varrho_{\max}(\mathcal{G}; n, k)$, then

- (1) for $n = 1$, we have $k = 0$ and $G \cong K_1$;
- (2) for $n = 2$, we have $k = 2$ and $G \cong K_2$;
- (3) for $n = 3$, we have $k = 0$ and $G \cong K_3$, or we have $k = 2$ and $G \cong P_3$;
- (4) for $n \geq 3$, $k = n - 1$, we have $G \cong S_n$;
- (5) for $n \geq 4$, $k = 1$, we have $G \cong P'_n$ where P'_n is as in Lemma 3.8;
- (6) for $n \geq 4$, $k = 2$, we have $G \cong P_n$;
- (7) for $n \geq 5$, $k = 3$, we have $G \cong D(n; 1, 2)$.

Proof. (1), (2), (3), (4) are obvious. (6), (7) follows from Theorem 3.7.

For a graph $H \in \mathcal{G}(n; 1)$, H is not a tree. Thus H contains cycles. If H is not a unicyclic graph. By delete some edges of H , we can get a connected unicyclic graph \mathcal{B} . By Lemma 3.1, then $\varrho(\mathcal{B}) > \varrho(H)$. By Lemma 3.8, we know that $\varrho(P'_n) \leq \varrho(\mathcal{B})$, with equality if and only if $\mathcal{B} \cong P'_n$. Note that there is only one pendant vertex in P'_n . Then (5) follows. \square

4. Conjecture

Based on computational evidence, we make the following conjecture.

Conjecture 4.1. If graph $G \in \mathcal{G}(n; k)$ ($4 \leq k \leq n - 2$) satisfying $\varrho(G) = \varrho_{\max}(\mathcal{G}; n, k)$, then $G \cong D(n; \lceil \frac{k}{2} \rceil, \lfloor \frac{k}{2} \rfloor)$.

Acknowledgments

We express many thanks to the referees for their kind reviews and helpful suggestions. The first author was supported by NSFC (Nos. 11171290, 11101057) and the fourth author was supported by NSFC (No. 11071078, No. 11075057).

References

- [1] R.L. Graham, H.O. Pollak, On the addressing problem for loop switching, Bell Syst. Tech. J. 50 (1971) 2495–2519.
- [2] M. Edelberg, M.R. Garey, R.L. Graham, On the distance matrix of a tree, Discrete Math. 14 (1976) 23–29.
- [3] R.L. Graham, L. Lovasz, Distance matrix polynomials of trees, Adv. Math. 29 (1978) 60–88.
- [4] R.L. Graham, H.O. Pollak, On embedding graphs in squashed cubes, in: Graph Theory and Applications, in: Lecture Notes in Mathematics, vol. 303, Springer, Berlin, 1972, pp. 99–110.
- [5] H. Hosoya, M. Murakami, M. Gotoh, Distance polynomial and characterization of a graph, Natur. Sci. Rep. Ochanomizu Univ. 24 (1973) 27–34.
- [6] D.H. Rouvray, The search for useful topological indices in chemistry, Am. Sci. 61 (1973) 729–735.
- [7] E.W. Dijkstra, A note on two problems in connection with graphs, Numer. Math. 1 (1959) 269–271.
- [8] M.L. Fredman, New bounds on the complexity of the shortest path problem, SIAM J. Comput. 5 (1976) 83–89.
- [9] O. Perron, Zur theorie der matrizen, Math. Ann. 64 (1907) 248–263.
- [10] B.F. Wu, E.L. Xiao, Y. Hong, The spectral radius of trees on k pendant vertices, Linear Algebra Appl. 395 (2005) 343–349.
- [11] Y. Hong, X.D. Zhang, Sharp upper and lower bounds for largest eigenvalue of the Laplacian matrices of trees, Disc. Math. 296 (2005) 187–197.
- [12] J.L. Shu, Y. Hong, R.K. Wen, A sharp upper bound on the largest eigenvalue of the Laplacian matrix of a graph, Linear Algebra Appl. 347 (2002) 123–129.
- [13] H. Minc, Nonnegative Matrices, John Wiley & Sons Inc., New York, 1988.
- [14] G.L. Yu, Y.R. Wu, J.L. Shu, Some graft transformations and its application on a distance spectrum, Disc. Math. 311 (2011) 2117–2123.
- [15] D. Stevanović, A. Ilić, Distance spectral radius of trees with fixed maximum degree, Electron. J. Linear Algebra 20 (2010) 168–179.
- [16] A. Ilić, Distance spectral radius of trees with given matching number, Discrete Appl. Math. 158 (2010) 1799–1806.